17:5		AND MEDICATION ORDER FOR SCHOOL AND CHILD CARE SETTINGS	
PARENT/GUARDIAN COMPLETE AND SIGN:			
Child Name:			
Parent/Guardian Name:			
Healthcare Provider Name:		Phone:	
Triggers: □Weather (cold air, wind) □Illness □Exercise □Smoke □Dust □Pollen □Other:			
Life threatening allergy, specify:			
youth presc relief	n, and if necessary, contact our hear ribed medication and supplies, an	to share this information, follow this plan, administer medication and care for my chilealthcare provider. I assume full responsibility for providing the school/program and to comply with board policies, if applicable. I am aware 911 may be called if a quichild/youth is experiencing symptoms. I approve this care plan for my child/youth. DATE NURSE/CCHC SIGNATURE DATE	
Distriction and the second sec			5-161.cm(n
HEALTHCARE PROVIDER COMPLETE ALL ITEMS,		QUICK RELIEF (RESCUE) MEDICATION: ☐ Albuterol ☐ Other: Common side effects: ↑ heart rate, tremor ☐ Have child use spacer with inhaler.	
SIGN AND DATE:		Controller medication used at home:	5.0
IF YOU SEE THIS:		DO THIS:	
GREEN ZONE: No Symptoms Pretreat	No current symptoms Doing usual activities	Pretreat strenuous activity: Not required Routine Student/Parent required Civa OUICK RELIEF MED 10.15 minutes before activity:	uest
EEN ZON Sympton Pretreat	Doing usual activities	Give QUICK RELIEF MED 10-15 minutes before activity: ☐ 2 puffs ☐ 4 puffs ☐ Repeat in 4 hours, if needed for additional physical activity.	
Syr Pre		If child is currently experiencing symptoms, follow YELLOW ZONE.	
R S		if china is currently experiencing symptoms, Johow Telegov Zowe.	
	Trouble breathing	1. Stop physical activity.	
ms eigh	Wheezing	2. Give QUICK RELIEF MED: ☐ 2 puffs ☐ 4 puffs	
207	Frequent cough	3. Stay with child/youth and maintain sitting position.	
3 (4)	Complains of tight chest	4. REPEAT QUICK RELIEF MED, if not improving in 15 minutes: 2 puffs 4 p	puffs
YELLOW ZONE: Mild symptoms	Not able to do activities,	5. Child/youth may go back to normal activities, once symptoms are relieved.	
画	but talking in complete	6. Notify parents/guardians and school nurse.	
	sentences	If symptoms do not improve or worsen, follow RED ZONE.	
MORE BUILDING	• Peak flow:&	1. Give QUICK RELIEF MED: ☐ 2 puffs ☐ 4 puffs	
5 "	Coughs constantlyStruggles to breathe	Refer to anaphylaxis plan, if child/youth has life-threatening allergy.	
GE GE	Trouble talking (only speaks)	2. Call 911 and inform EMS the reason for the call.	
pto	3-5 words)	3. Stay with child/youth. Remain calm, encouraging slower, deeper breaths.	
Z E	• Skin of chest and/or neck	4. Notify parents/guardians and school nurse.	
RED ZONE: EMERGENC Severe Symptoms	pull in with breathing	5. If symptoms do not improve, REPEAT QUICK RELIEF MED: 2 puffs 4 put	ffs
ON Je	• Lips/fingernails gray or blue	every 5 minutes until EMS arrives.	
D Z Se	 ◆ Level of consciousness 	School personnel should not drive student to hospital.	
핆	• Peak flow <		
PROVIDER INSTRUCTIONS FOR QUICK RELIEF INHALER USE: CHECK APPROPRIATE BOX(ES)			
Student needs supervision or assistance to use inhaler. Student will not self-carry inhaler.			
Student understands proper use of asthma medications, and in my opinion, can carry and use his/her inhaler at school			
independently with approval from school nurse and completion of contract.			
Student will notify school staff after using quick relief inhaler, if symptoms do not improve with use.			
HEALTH (CARE PROVIDER SIGNATURE	PRINT PROVIDER NAME DATE FAX PHONE	
Copies	of plan provided to: \square Teacher(s)		

